

## CCL18 blocking peptide

---

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-CCL18 Antibody (Catalog no. 4076257) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5076257
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	CCL18
UniProt ID	P55774
Purity	>85%
Synonyms	AMAC1; DCCK1; MIP4; PARC; SCYA18; C-C motif chemokine 18; Alternative macrophage activation-associated CC chemokine 1; AMAC-1; CC chemokine PARC; Dendritic cell chemokine 1; DC-CK1; Macrophage inflammatory protein 4; MIP-4; Pulmonary and activation-regulated chemokine; Small-inducible cytokine A18

Storage/Stability	-20°C/1 year
Shipping	Gel Packs

[www.realgenelabs.com](http://www.realgenelabs.com)

For Research Use only