

Ig lambda C blocking peptide

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-Ig lambda C Antibody (Catalog no. 4074436) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5074436
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	IGLC1; IGLC2; IGLC3; IGLC6; IGLC7
UniProt ID	P15814; P0CG04; P0CG05; P0CG06; P0CF74; A0M8Q6
Purity	>85%

Synonyms	IGLL1; IGL1; Immunoglobulin lambda-like polypeptide 1; CD179 antigen-like family member B; Ig lambda-5; Immunoglobulin omega polypeptide; Immunoglobulin-related protein 14.1; CD179b; IGLC1; Ig lambda-1 chain C regions; IGLC2; Ig lambda-2 chain C regions; IGLC3; Ig lambda-3 chain C regions; IGLC6; Ig lambda-6 chain C region; IGLC7; Ig lambda-7 chain C region
Storage/Stability	-20°C/1 year
Shipping	Gel Packs

www.realgenelabs.com

For Research Use only