

## NMDAR2D blocking peptide

---

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-NMDAR2D Antibody (Catalog no. 4074423) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5074423
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	GRIN2D
UniProt ID	O15399
Purity	>85%
Synonyms	GluN2D; NMDAR2D; Glutamate receptor ionotropic, NMDA 2D; GluN2D; EB11; Glutamate [NMDA] receptor subunit epsilon-4; N-methyl D-aspartate receptor subtype 2D; NMDAR2D; NR2D
Storage/Stability	-20°C/1 year
Shipping	Gel Packs

