

PI3K p110 gamma blocking peptide

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-PI3K p110 gamma Antibody (Catalog no. 4075034) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5075034
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	PIK3CG
UniProt ID	P48736
Purity	>85%

Synonyms	Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit gamma isoform; PI3-kinase subunit gamma; PI3K-gamma; PI3Kgamma; PtdIns-3-kinase subunit gamma; Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit gamma; PtdIns-3-kinase subunit p110-gamma; p110gamma; Phosphoinositide-3-kinase catalytic gamma polypeptide; Serine/threonine protein kinase PIK3CG; p120-PI3K
Storage/Stability	-20°C/1 year
Shipping	Gel Packs

www.realgenelabs.com

For Research Use only