

## PLC gamma 1 (pY783) blocking peptide

---

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-PLC gamma 1 (pY783) Antibody (Catalog no. 4073290) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5173290
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	PLCG1
UniProt ID	P19174
Purity	>85%
Synonyms	PLC1; 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1; PLC-148; Phosphoinositide phospholipase C-gamma-1; Phospholipase C-II; PLC-II; Phospholipase C-gamma-1; PLC-gamma-1
Storage/Stability	-20°C/1 year

Shipping

Gel Packs

[www.realgenelabs.com](http://www.realgenelabs.com)

For Research Use only