

## RAC1/2/3 (pS71) blocking peptide

---

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-RAC1/2/3 (pS71) Antibody (Catalog no. 4076039) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5176039
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	RAC1; RAC2; RAC3
UniProt ID	P63000; P15153; P60763
Purity	>85%

Synonyms	RAC1; TC25; MIG5; Ras-related C3 botulinum toxin substrate 1; Cell migration-inducing gene 5 protein; Ras-like protein TC25; p21-Rac1; RAC2; Ras-related C3 botulinum toxin substrate 2; GX; Small G protein; p21-Rac2; RAC3; Ras-related C3 botulinum toxin substrate 3; p21-Rac3; CDC42; Cell division control protein 42 homolog; G25K GTP-binding protein
Storage/Stability	-20°C/1 year
Shipping	Gel Packs

[www.realgenelabs.com](http://www.realgenelabs.com)

For Research Use only