

TAO1 blocking peptide

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunobloting) when non-specific binding is an issue.

This peptide is recommended to block anti-TAO1 Antibody (Catalog no. 4074843) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

| Catalog No. | 5074843 |
|------------------|--|
| Size | lmg |
| Product Category | Blocking Peptide |
| Gene Symbol | TAOK1 |
| UniProt ID | Q7L7X3 |
| Purity | >85% |
| Synonyms | KIAA1361; MAP3K16; MARKK; Serine/threonine-protein kinase TAO1; Kinase from chicken homolog B; hKFC-B; MARK Kinase; MARKK; Prostate-derived sterile 20-like kinase 2; PSK-2; PSK2; Prostate-derived STE20-like kinase 2; Thousand and one amino acid protein kinase 1; TAOK1; hTAOK1 |

| Storage/Stability | -20°C/1 year |
|-------------------|--------------|
| Shipping | Gel Packs |

www.realgenelabs.com

For Research Use only