

4x Stacking Gel Separatng Buffer [pH 8.8]

PAGE Separation Buffer [pH 8.8] is used in the preparation of separating or resolving gel mix while performing SDS-PAGE which is used for the separation of proteins through electrophoresis and it is based on the fact that charged molecules will migrate through a matrix upon application of an electrical field. The matrix for protein electrophoresis separation is polyacrylamide. SDS-PAGE uses two types of buffer systems: the continuous buffer system and the discontinuous buffer system. In the continuous buffer system, the pH of the gel matrix remains constant throughout the separation. In contrast, the discontinuous buffer system consists of a narrow layer of stacking gel (of large pore size and acidic pH) above the main separating or resolving gel matrix of alkaline pH (pH 8.8). The stacking gel concentrates the protein sample before entering the separating gel and hence enhancing resolution. SDS-PAGE with a discontinuous buffer system is the most popular electrophoresis technique used to analyze polypeptides.

The buffer consist Tris Base & SDS in recommended concentrations at pH 8.8. The buffer is supplied in 4x contcentrate and should be diluted prior to use in electrophoresis.

Catalog No.	503022
Size	100mL / 500mL / Size1 / WSP / Size2 / WSP
Product Category	SDS-PAGE Buffer & Stain
Storage/Stability	Ambient/1 year
Shipping	Ambient